

Precautions for Individuals Taking Blood Thinning Medications

As certain blood thinners can make it difficult for a wound to stop bleeding, please watch out for any signs of uncontrollable bleeding after the injection. Please also let us know which medication you are taking on the Prevacination Screening Questionnaire.

Precautions about the intramuscular injection



Please apply pressure on the injection site for at least 2 minutes after the injection.

- If your arm swells up, feels numb or has any other symptoms, please consult a doctor.
- There is no need to stop taking your medication prior to this injection.

Please check if you are taking any of the anticoagulant or antiplatelet medicines below.

Anticoagulant Medicine

The following medicines are usually administered after surgery for the heart, arrhythmia or thrombosis.

Brand-name	Generic Name
Warfarin	Warfarin potassium
Pradaxa	Dabigatran
Xarelto	Rivaroxaban
Eliquis	Apixaban
Lixiana	Edoxaban tosilate hydrate

Antiplatelet Medicine

The following medicines are commonly administered for arteriosclerosis, angina, myocardial infarction, cerebral infarction (stroke) and arteriosclerosis obliterans.

Brand-name	Generic Name
Bayaspirin	Aspirin
Panaldine	Ticlopidine
Plavix	Clopidogrel
Efient	Prasugel
Bufferin	Aspirin/ dialuminate
Brilinta	Ticagrelor

Brand-name	Generic Name
ConPlavin	Aspirin and clopidogrel
Takelda	Aspirin and lansoprazole
Epadel	Ethyl icosapentate
Dorner, Procylin	Beraprost sodium
Anplag	Sarpogrelate hydrochloride